ProbStat Forum, Volume 08, July 2015, Pages 103–111 ISSN 0974-3235

Some characterization results based on conditional expectation of dual generalized order statistics

Haseeb Athar, Zuber Akhter^{\dagger}

Department of Statistics and Operations Research, Aligarh Muslim University, Aligarh-202 002. India.

Abstract. In this paper, a general form of continuous probability distributions $F(x) = a(h(x))^{-c}$ $x \in (\alpha, \beta)$ has been characterized through conditional expectation of p - th power of difference of two dual generalized order statistics. Further, some deductions and related results are also discussed.

1. Introduction

Kamps (1995) introduced the concept of generalized order statistics (gos) to unify several models of ascendingly ordered random variables, e.g. upper order statistics, k-record values, progressively Type-II censored order statistics, Pfeifer records and sequential order statistics. These models can be effectively applied in reliability theory and survival analysis. However, random variables that are decreasingly ordered cannot be comprised into this framework. Further, this model is inappropriate to study, e.g. reversed ordered order statistic and lower record values models. Based on the gos, Burkschat $et \ al.$ (2003) introduced the concept of the dual generalized order statistics (dgos) that enables a common approach to study descendingly ordered random variables like reversed ordered order statistics and lower record values.

Let $n \ge 2$ be a given integer and $\tilde{m} = (m_1, m_2, \dots, m_{n-1}) \in \Re^{n-1}, k > 0$ be the parameters such that

$$\gamma_i = k + n - i + \sum_{j=i}^{n-1} m_j > 0 \text{ for } 1 \le i \le n-1$$

The random variables $X_d(1, n, \tilde{m}, k), X_d(2, n, \tilde{m}, k), \ldots, X_d(n, n, \tilde{m}, k)$ are said to be dual generalized order statistics from an absolutely continuous distribution function df F() with the probability density function pdf f(), if their joint density function is of the form

$$f_{X_d(1,n,\tilde{m},k),\ldots,X_d(n,n,\tilde{m},k)}(x_1,x_2,\ldots,x_n)$$

$$= k \left(\prod_{j=1}^{n-1} \gamma_j\right) \left(\prod_{i=1}^{n-1} \left[F(x_i)\right]^{m_i} f(x_i)\right) \left[F(x_n)\right]^{k-1} f(x_n)$$

$$\tag{1}$$

²⁰¹⁰ Mathematics Subject Classification. Primary 62E10, 62G30

Keywords. Dual generalized order statistics, lower order statistics, record values, conditional expectation, characterization. Received: 11 September 2014; Accepted: 01 July 2015.

[†]Corresponding author

Email address: akhterzuber022@gmail.com (Zuber Akhter[†])

104

for $F^{-1}(1) > x_1 \ge x_2 \ge \ldots \ge x_n > F^{-1}(0)$.

Here we may consider two cases:

Case I: $\gamma_i \neq \gamma_j; i \neq j = 1, 2, ..., n - 1.$

In view of (1) the pdf of r - th dual generalized order statistics $X_d(r, n, \tilde{m}, k)$ is

$$f_{X_d(r,n,\tilde{m},k)}(x) = C_{r-1}f(x)\sum_{i=1}^r a_i(r) [F(x)]^{\gamma_i - 1}$$
(2)

and the joint $pd\!f$ of $X_d(r,n,\tilde{m},k)$ and $X_d(s,n,\tilde{m},k),\, 1 \leq r < s \leq n$ is

$$f_{X_d(r,n,\tilde{m},k),X_d(s,n,\tilde{m},k)}(x,y) = C_{s-1} \left(\sum_{i=r+1}^s a_i^{(r)}(s) \left[\frac{F(y)}{F(x)} \right]^{\gamma_i} \right)$$
$$\times \left(\sum_{i=1}^r a_i(r) \left[F(x) \right]^{\gamma_i} \right) \frac{f(x)}{F(x)} \frac{f(y)}{F(y)}, \ \alpha \le y < x \le \beta.$$
(3)

The conditional pdf of $X_d(s, n, \tilde{m}, k)$ given $X_d(r, n, \tilde{m}, k) = x, 1 \le r < s \le n$ in view of (2) and (3) is

$$f_{X_d(s,n,\tilde{m},k)|X_d(r,n,\tilde{m},k)}(y|x) = \frac{C_{s-1}}{C_{r-1}} \sum_{i=r+1}^s a_i^{(r)}(s) \frac{[F(y)]^{\gamma_i - 1}}{[F(x)]^{\gamma_i}} f(y), \ x > y.$$

$$\tag{4}$$

where,

$$C_{r-1} = \prod_{i=1}^{r} \gamma_i, \ \gamma_i = k + n - i + \sum_{j=i}^{n-1} m_j > 0,$$
$$a_i(r) = \prod_{\substack{j=1\\ j \neq i}}^{r} \frac{1}{(\gamma_j - \gamma_i)}, \ \gamma_i \neq \gamma_j, \ 1 \le i \le r \le n$$

and

$$a_i^{(r)}(s) = \prod_{\substack{j=r+1\\j\neq i}}^s \frac{1}{(\gamma_j - \gamma_i)}, \ \gamma_i \neq \gamma_j, \ r+1 \le i \le s \le n.$$

Case II: $m_i = m$ (say); i = 1, 2, ..., n - 1.

The *pdf* of r - th of $X_d(r, n, m, k)$ is

$$f_{X_d(r,n,m,k)}(x) = \frac{C_{r-1}}{(r-1)!} \left[F(x) \right]^{\gamma_r - 1} f(x) \ g_m^{r-1} \left(F(x) \right)$$
(5)

and the joint pdf of $X_d(r, n, m, k)$ and $X_d(s, n, m, k), 1 \le r < s \le n$ is

$$f_{X_d(r,n,m,k),X_d(s,n,m,k)}(x,y) = \frac{C_{s-1}}{(r-1)!(s-r-1)!} [F(x)]^m f(x) g_m^{r-1}(F(x)) \\ \times [h_m(F(y)) - h_m(F(x))]^{s-r-1} [F(y)]^{\gamma_s - 1} f(y), \ \alpha \le y < x \le \beta.$$
(6)

The conditional pdf of $X_d(s, n, m, k)$ given $X_d(r, n, m, k) = x, 1 \le r < s \le n$ in view of (5) and (6) is

$$f_{X_d(s,n,m,k)|X_d(r,n,m,k)}(y|x) = \frac{C_{s-1}}{C_{r-1}(s-r-1)!(m+1)^{s-r-1}} \frac{[F(y)]^{\gamma_{s-1}}}{[F(x)]^{\gamma_{r+1}}} \times \left[[F(x)]^{m+1} - [F(y)]^{m+1} \right]^{s-r-1} f(y), \ x > y.$$
(7)

where,

$$h_m(x) = \begin{cases} -\frac{1}{m+1} x^{m+1} & , \ m \neq -1 \\ -\log x & , \ m = -1 \end{cases}$$

and
$$g_m(x) = h_m(x) - h_m(1), x \in [0, 1).$$

If m = 0 and k = 1, then $X_d(r, n, m, k)$ reduces to the (n - r + 1) - th lower order statistic, $X_{n-r+1:n}$ from the sample X_1, X_2, \ldots, X_n [David and Nagaraja (2003)]. If m = -1 and k = 1, then $X_d(r, n, m, k)$ is the r - th lower record value from an infinite sequence of independent and identically distributed *(iid)* random variables (rv) [Ahsanullah (1995)].

The characterization of probability distributions through conditional expectation of order statistics and record values have been seen among others are Nagraja (1977, 1988), Wu and Ouyang (1996), Franco and Ruiz (1995, 1996, 1997), López-Blázquez and Moreno-Rebello (1997), Wesolowski and Ahsanullah (1997), Dembińska and Wesolowski (1998, 2000), Wu and Lee (2001), Raqab (2002), Lee (2003), Athar *et al.* (2003), Khan and Athar (2004), Wu (2004), Noor and Athar (2014), Athar and Akhter (2015) and references therein.

Khan and Abu-Salih (1989) characterized some general forms of distributions through conditional expectation of order statistics fixing adjacent order statistics. Khan and Abouanmoh (2000) extended the result of Khan and Abu-Salih (1989) and characterized the distributions when the conditioning is not adjacent. Khan *et al.* (2006) established characterizing relationships for the distributions through *gos* and characterized several distributions through conditional expectation of function of *gos*. Further, Khan *et al.* (2010) characterized several distributions through conditional expectation of function of *dgos*.

For various developments on characterization dealing with gos and dgos one may refer to Keseling (1999), Bieniek and Szynal (2003), Ahsanullah (2004), Khan and Alzaid (2004), Mbah and Ahsanullah (2007), Bieniek (2007, 2009), Samuel (2008), Khan et al. (2012), Noor et al. (2014, 2015) and references therein.

2. Characterization theorems

Theorem 2.1: Let $X_d(i, n, \tilde{m}, k)$, $1 \leq i \leq n$, be dual generalized order statistics based on absolutely continuous df F(x) and pdf f(x) over the support (α, β) , where α and β may be finite or infinite, then for $1 \leq r < s \leq n$,

$$E\left[\{h(X_d(s, n, \tilde{m}, k)) - h(X_d(r, n, \tilde{m}, k))\}^p | X_d(r, n, \tilde{m}, k) = x\right] = \xi_{r,s,p}(x)$$
$$= a^*(h(x))^p \tag{8}$$

if and only if

$$F(x) = a(h(x))^{-c}, \ a \neq 0,$$
(9)

where

$$a^* = \sum_{j=0}^p (-1)^{j+p} \binom{p}{j} \prod_{i=r+1}^s \binom{c\gamma_i}{c\gamma_i - j}, \ c\gamma_i \neq j$$

and h(x) is a monotonic and differentiable function of x and p is a positive integer.

Proof: To prove the necessary part, we have

$$E\left[\{h(X_d(s, n, \tilde{m}, k)) - h(X_d(r, n, \tilde{m}, k))\}^p | X_d(r, n, \tilde{m}, k) = x\right]$$

= $\frac{C_{s-1}}{C_{r-1}} \int_{\alpha}^{x} (h(y) - h(x))^p \sum_{i=r+1}^{s} a_i^{(r)}(s) \left[\frac{F(y)}{F(x)}\right]^{\gamma_i - 1} \frac{f(y)}{F(x)} dy.$ (10)

Let

$$t = \left[\frac{F(y)}{F(x)}\right]$$
, which implies $(h(y) - h(x))^p = (-1)^p (h(x))^p \left(1 - t^{-1/c}\right)^p$.

Then the right hand side of (10) reduces to

$$E\left[\left\{h(X_d(s, n, \tilde{m}, k)) - h(X_d(r, n, \tilde{m}, k))\right\}^p | X_d(r, n, \tilde{m}, k) = x\right]$$

$$= \frac{C_{s-1}}{C_{r-1}} \sum_{i=r+1}^s a_i^{(r)}(s) \int_0^1 (-1)^p (h(x))^p (1 - t^{-1/c})^p t^{\gamma_i - 1} dt$$

$$= \frac{C_{s-1}}{C_{r-1}} (h(x))^p \sum_{i=r+1}^s a_i^{(r)}(s) \left(\sum_{j=0}^p (-1)^{j+p} \binom{p}{j} \int_0^1 t^{\gamma_i - \frac{j}{c} - 1} dt\right)$$

$$= (h(x))^p \sum_{j=0}^p (-1)^{j+p} \binom{p}{j} \left(\frac{C_{s-1}}{C_{r-1}} \sum_{i=r+1}^s a_i^{(r)}(s) \left(\frac{c}{c\gamma_i - j}\right)\right).$$

This proves the necessary part.

To prove sufficiency part, let

$$E\left[\{h(X_d(s, n, \tilde{m}, k)) - h(X_d(r, n, \tilde{m}, k))\}^p | X_d(r, n, \tilde{m}, k) = x\right] = \xi_{r,s,p}(x).$$

Therefore,

$$\frac{C_{s-1}}{C_{r-1}} \int_{\alpha}^{x} (h(y) - h(x))^{p} \sum_{i=r+1}^{s} a_{i}^{(r)}(s) \frac{[F(y)]^{\gamma_{i}-1}}{[F(x)]^{\gamma_{i}}} f(y) dy = \xi_{r,s,p}(x).$$

Differentiating both sides w.r.t. x, we get

$$-ph'(x)\frac{C_{s-1}}{C_{r-1}}\int_{\alpha}^{x}(h(y)-h(x))^{p-1}\sum_{i=r+1}^{s}a_{i}^{(r)}(s)\frac{[F(y)]^{\gamma_{i}-1}}{[F(x)]^{\gamma_{i}}}f(y)dy$$
$$-\frac{C_{s-1}}{C_{r-1}}\int_{\alpha}^{x}(h(y)-h(x))^{p}\sum_{i=r+1}^{s}\gamma_{i}\ a_{i}^{(r)}(s)\frac{[F(y)]^{\gamma_{i}-1}}{[F(x)]^{\gamma_{i}}}\frac{f(x)}{F(x)}f(y)dy = \xi_{r,s,p}'(x),$$

or,

$$-ph'(x)\xi_{r,s,p-1}(x) - \frac{C_{s-1}}{C_{r-1}}\int_{\alpha}^{x} (h(y) - h(x))^{p} \sum_{i=r+1}^{s} \gamma_{i} \ a_{i}^{(r)}(s) \frac{[F(y)]^{\gamma_{i}-1}}{[F(x)]^{\gamma_{i}}} \frac{f(x)}{F(x)} f(y)dy = \xi_{r,s,p}'(x).$$

Since,

$$a_i^{(r+1)}(s) = (\gamma_{r+1} - \gamma_i)a_i^{(r)}(s)$$
 and $C_r = \gamma_{r+1}C_{r-1}$.

Therefore, we have

$$-\gamma_{r+1} \frac{C_{s-1}}{C_{r-1}} \frac{f(x)}{F(x)} \int_{\alpha}^{x} (h(y) - h(x))^{p} \sum_{i=r+1}^{s} a_{i}^{(r)}(s) \frac{[F(y)]^{\gamma_{i}-1}}{[F(x)]^{\gamma_{i}}} f(y) dy$$

+ $\gamma_{r+1} \frac{C_{s-1}}{C_{r}} \frac{f(x)}{F(x)} \int_{\alpha}^{x} (h(y) - h(x))^{p} \sum_{i=r+2}^{s} a_{i}^{(r+1)}(s) \frac{[F(y)]^{\gamma_{i}-1}}{[F(x)]^{\gamma_{i}}} f(y) dy$
= $\xi_{r,s,p}'(x) + ph'(x)\xi_{r,s,p-1}(x).$ (11)

Rearranging the terms of (11), we get

$$\frac{f(x)}{F(x)} = -\frac{1}{\gamma_{r+1}} \frac{\xi'_{r,s,p}(x) + ph'(x)\xi_{r,s,p-1}(x)}{[\xi_{r,s,p}(x) - \xi_{r+1,s,p}(x)]}.$$
(12)

Now consider

$$\begin{aligned} \xi'_{r,s,p}(x) + p \, h'(x) \xi_{r,s,p-1}(x) \\ &= p \, h'(x)(h(x))^{p-1} \sum_{j=0}^{p} (-1)^{j+p} {p \choose j} \prod_{i=r+1}^{s} \left(\frac{c\gamma_{i}}{c\gamma_{i} - j} \right) \\ &+ p \, h'(x)(h(x))^{p-1} \sum_{j=0}^{p-1} (-1)^{j+p-1} {p-1 \choose j} \prod_{i=r+1}^{s} \left(\frac{c\gamma_{i}}{c\gamma_{i} - j} \right) \\ &= p \, h'(x)(h(x))^{p-1} \left[\sum_{j=0}^{p} (-1)^{j+p} {p \choose j} \prod_{i=r+1}^{s} \left(\frac{c\gamma_{i}}{c\gamma_{i} - j} \right) \\ &- \sum_{j=0}^{p-1} (-1)^{j+p} {p \choose j} \frac{(p-j)}{p} \prod_{i=r+1}^{s} \left(\frac{c\gamma_{i}}{c\gamma_{i} - j} \right) \right] \\ &= h'(x)(h(x))^{p-1} \left[\sum_{j=0}^{p} (-1)^{j+p} {p \choose j} j \prod_{i=r+1}^{s} \left(\frac{c\gamma_{i}}{c\gamma_{i} - j} \right) \right] \end{aligned}$$
(13)

 $\quad \text{and} \quad$

$$\begin{aligned} \xi_{r,s,p}(x) &- \xi_{r+1,s,p}(x) \\ &= (h(x))^p \left[\sum_{j=0}^p (-1)^{j+p} \binom{p}{j} \prod_{i=r+1}^s \binom{c\gamma_i}{c\gamma_i - j} - \sum_{j=0}^p (-1)^{j+p} \binom{p}{j} \prod_{i=r+2}^s \binom{c\gamma_i}{c\gamma_i - j} \right] \\ &= (h(x))^p \left[\sum_{j=0}^p (-1)^{j+p} \binom{p}{j} \prod_{i=r+2}^s \binom{c\gamma_i}{c\gamma_i - j} \binom{c\gamma_{r+1}}{c\gamma_{r+1} - j} - 1 \right] \end{aligned}$$

Athar and Akthar / ProbStat Forum, Volume 08, July 2015, Pages 103-111

$$= (h(x))^p \left[\frac{1}{c\gamma_{r+1}} \sum_{j=0}^p (-1)^{j+p} \binom{p}{j} j \prod_{i=r+1}^s \binom{c\gamma_i}{c\gamma_i - j} \right].$$

$$\tag{14}$$

Therefore in view of (12), we get

$$\frac{f(x)}{F(x)} = -\frac{1}{\gamma_{r+1}} \frac{h'(x)(h(x))^{p-1}}{(h(x))^p} \frac{\left[\sum_{j=0}^p (-1)^{j+p} \binom{p}{j} j \prod_{i=r+1}^s \binom{c\gamma_i}{c\gamma_i - j}\right]}{\left[\frac{1}{c\gamma_{r+1}} \sum_{j=0}^p (-1)^{j+p} \binom{p}{j} j \prod_{i=r+1}^s \binom{c\gamma_i}{c\gamma_i - j}\right]}$$
$$= -\frac{c h'(x)}{h(x)}.$$

Implying that

$$F(x) = a(h(x))^{-c}.$$

Similarly, the characterization result for case II may be obtained on the lines of Theorem 2.1. Corollary 2.1: Under the condition as stated in Theorem 2.1,

$$E[h(X_d(s, n, \tilde{m}, k))|X_d(r, n, \tilde{m}, k) = x] = h(x) \prod_{i=r+1}^s \left(\frac{c\gamma_i}{c\gamma_i - 1}\right), \ c\gamma_i \neq 1,$$
(15)

and consequently

$$E[h(X_d(r+1, n, \tilde{m}, k))|X_d(r, n, \tilde{m}, k) = x] = h(x) \left(\frac{c\gamma_{r+1}}{c\gamma_{r+1} - 1}\right), \ c\gamma_{r+1} \neq 1$$
(16)

if and only if

$$F(x) = a(h(x))^{-c}, \quad a \neq 0,$$

where h(x) is a monotonic and differentiable function of x.

Proof: Expression (15) can be proved in view of Theorem 2.1 at p = 1 and (16) can be obtained at s = r + 1 in (15).

Remark 2.1: Let $m_i = m = 0$, i = 1, 2, ..., n - 1 and k = 1, then, characterization result for lower order statistics is given as

$$E[\{h(X_{n-s+1:n}) - h(X_{n-r+1:n})\}^p | X_{n-r+1:n} = x] = b^*(h(x))^p$$

if and only if

$$F(x) = a(h(x))^{-c}, \ a \neq 0,$$

where

$$b^* = \sum_{j=0}^{p} (-1)^{j+p} \binom{p}{j} \prod_{i=r+1}^{s} \binom{c(n-i+1)}{c(n-i+1)-j}, \ c(n-i+1) \neq j.$$

Remark 2.2: At m = -1 and k = 1, the characterization result for lower record will be

$$E[\{h(X_{L(s)}) - h(X_{L(r)})\}^p | X_{L(r)} = x] = c^*(h(x))^p$$

if and only if

$$F(x) = a(h(x))^{-c}, \ a \neq 0,$$

where

$$c^* = \sum_{i=0}^{p} (-1)^{i+p} {p \choose i} \left(\frac{c}{c-i}\right)^{s-r}, \ c \neq i.$$

3. Examples

(i) Power function distribution

$$F(x) = x^{\nu}, \quad 0 < x \le 1, \ \nu > 0.$$

Then F(x) is given by (9) with a = 1, $c = -\nu$ and h(x) = x,

(ii) Inverse Power distribution

$$F(x) = \left(\frac{x-\alpha}{\beta-\alpha}\right)^{\theta}, \ \alpha < x < \beta, \ \theta > 0, \ \alpha, \beta \in \mathbb{R}, \ \alpha < \beta.$$

Then F(x) is given by (9) with a = 1, $c = -\theta$ and $h(x) = \left(\frac{x - \alpha}{\beta - \alpha}\right)$.

(iii) Reflected exponential distribution

$$F(x) = e^{\lambda(x-\mu)}, \quad -\infty < x < \mu, \ \lambda > 0.$$

Then F(x) is given by (9) with a = 1, c = -1 and $h(x) = e^{\lambda(x-\mu)}$.

(iv) Inverse Weibull distribution

$$F(x) = e^{-\theta x^{-\nu}}, \quad 0 < x < \infty, \ \theta, \nu > 0.$$

Then F(x) is given by (9) with a = 1, $c = \theta$ and $h(x) = e^{x^{-\nu}}$.

(v) Inverse exponential distribution

$$F(x) = e^{-\frac{\theta}{x}}, \quad x > 0, \ \theta > 0.$$

· - ·/

Then F(x) is given by (9) with a = 1, $c = \theta$ and $h(x) = e^{\frac{1}{x}}$.

(vi) Gumbel distribution

$$F(x) = e^{-e^{-x}}, \quad -\infty < x < \infty.$$

Then F(x) is given by (9) with a = 1, c = 1 and $h(x) = e^{e^{-x}}$.

(vii) Burr type X distribution

$$F(x) = (1 - e^{-x^2})^k, \quad 0 < x < \infty, \ k > 0.$$

Then F(x) is given by (9) with a = 1, $c = -\frac{k}{q}$ and $h(x) = (1 - e^{-x^2})^q$, $q \neq 0$.

Similarly, characterization results for other distributions may be obtained with proper choice of a and h(x). One may refer to Khan *et al.* (2007).

References

- [1] Ahsanullah, M. (1995): Record Statistics. Nova Science publishers, New York.
- [2] Ahsanullah, M. (2004): A characterization of the uniform distribution by dual generalized order statistics. Commun. Statist. Theor. Meth. 33, 2921-2928.
- [3] Athar, H., Yaqub, M. and Islam, H.M. (2003): On characterization of distributions through linear regression of record values and order statistics. *Aligarh J. Statist.*, 23, 97-105.
- [4] Athar, H. and Akhter, Z. (2015): Some characterization of continuous distributions based on order statistics. Int. J. Comp. Theo. Stat., 2(1), 31-36.
- [5] Bieniek, M. and Szynal, D. (2003): Characterizations of distributions via linearity of regression of generalized order statistics. *Metrika*, 58, 259-271.
- [6] Bieniek, M. (2007): On characterizations of distributions by regression of adjacent generalized order statistics. *Metrika*, 66, 233-242.
- [7] Bieniek, M. (2009): A note on characterizations of distributions by regressions of nonadjacent generalized order statistics. Aust. N. Z. J. Stat., 51, 89-99.
- [8] Burkschat, M., Cramer, E. and Kamps, U. (2003): Dual generalized order statistics. Metron, LXI(1), 13 26.
- [9] David, H.A. and Nagaraja, H.N. (2003): Order Statistics. John Wiley, New York.
- [10] Dembińska, A. and Wesolowski, J. (1998): Linearity of regression for non-adjacent order statistics. Metrika, 48, 215-222.
 [11] Dembińska, A. and Wesolowski, J. (2000): Linearity of regression for non-adjacent record values. J. Statist. Plann. Inference, 90, 195-205.
- [12] Franco, M. and Ruiz, J.M. (1995): On characterization of distributions with adjacent order statistics. *Statistics*, **26**, 375-385.
- [13] Franco, M. and Ruiz, J.M. (1996): On characterization of continuous distributions by conditional expectation of record values. Sankhyā Ser. A, 58, 135-141.
- [14] Franco, M. and Ruiz, J.M. (1997): On characterization of distributions by expected values of order statistics and record values with gap. *Metrika*, **45**, 107-119.
- [15] Kamps, U. (1995): A concept of generalized order statistics. J. Statist. Plann. Inference, 48, 1-23.
- [16] Keseling, C. (1999): Conditional distributions of generalized order statistics and some charactrizations. Metrika, 49, 27-40.
- [17] Khan, A.H. and Abu-Salih, M.S. (1989): Characterizations of probability distributions by conditional expectation of order statistics. *Metron*, 47, 171–181.
- [18] Khan, A.H. and Abouammoh (2000): Characterizations of distributions by conditional expectation of order statistics. J. Appl. Statist. Sci., 9, 159–167.
- [19] Khan, A.H. and Alzaid, A.A. (2004): Characterization of distributions through linear regression of non-adjacent generalized order statistics. J. Appl. Statist. Sci., 13, 123-136.
- [20] Khan, A.H. and Athar, H. (2004): Characterization of distributions through order statistics. J. Appl. Statist. Sci., 13(2), 147-154.
- [21] Khan, A.H., Khan, R. U. and Yaqub, M. (2006): Characterization of continuous distributions through conditional expectation of generalized order statistics. J. Appl. Prob. Statist., 1, 115-131.
- [22] Khan, A. H., Athar, H. and Chishti, S. (2007): On characterization of continuous distributions conditioned on a pair of order statistics. Journal of Applied Statistical Sciences, 16(3), 331-346.
- [23] Khan, A.H., Anwar, Z. and Chishti, S. (2010): Characterization of continuous distributions through conditional expectation of functions of dual generalized order statistics. Pak. J. Statist., 26(4), 615–628.
- [24] Khan, A.H., Shah, I.A. and Ahsanullah, M. (2012) Characterization through distributional properties of dual generalized order statistics, J. Egypt. Math. Soc., 20, 211214.
- [25] Lee, M.Y. (2003): Characterizations of the Pareto distribution by conditional expectations of record values. Commun. Korean Math. Soc., 18(1), 127-131.
- [26] López-Blázquez, F. and Moreno-Rebollo, J.L. (1997): A characterization of distributions based on linear regression of order statistics and record values. Sankhyā Ser. A, 59, 311-323.
- [27] Mbah, A.K. and. Ahsanullah, M. (2007): Some characterization of the power function distribution based on lower generalized order statistics. Pakistan J. Statist., 23, 139-146.
- [28] Nagaraja, H.N. (1977): On a characterization based on record values. Austral. J. Statist., 19(1), 70-73.
- [29] Nagaraja, H.N. (1988): Some characterizations of continuous distributions based on regressions of adjacent order statistics and record values. Sankhyā Ser. A, 50, 70-73.
- [30] Noor, Z. and Athar, H. (2014): Characterization of probability distributions by conditional expectation of function of record statistics. J. Egyptian Math. Soc., 22, 275-279.
- [31] Noor, Z., Athar, H. and Akhter, Z. (2014): Conditional expectation of generalized order statistics and characterization of probability distributions. J. Stat. Appl. Pro. Lett., 1(1), 9-18.
- [32] Noor, Z., Akhter, Z. and Athar, H. (2015): On characterization of probability distributions through conditional expectation of generalized and dual generalized order statistics. *Pak. J. Statist.*, **31(2)**, 159-170.
- [33] Raqab, M.Z. (2002): Inferences for generalized exponential distribution based on record statistics. J. Statist. Plann. Inference, 104(2), 339-350.
- [34] Samuel, P. (2008): Characterization of distributions by conditional expectation of generalized order statistics. *Statist. Papers*, **49**, 101-108.

- [35] Wesolowski, J. and Ahsanullah, M. (1997): On characterizing distributions via linearity of regression for order statistics. Aust. J. Stat., 39, 69–78.
- [36] Wu, J. W. and Ouyang, I.Y. (1996): On characterizing distributions by conditional expectations of the function of order statistics. *Metrika*, 43, 135–147.
- [37] Wu, J.W. and Lee, W.C. (2001): On characterization of generalized extreme value, power function, generalized Pareto and classical Pareto distributions by conditional expectation of record values. *Statist. Papers*, **42**, 225-242.
- [38] Wu, J.W. (2004): On characterizing distributions by conditional expectation of functions of non-adjacent record values. J. Appl. Statist. Sci., 13, 137-145.